

Simplicial Euclidean Relativistic Lagrangian

Marius I. Piso

Institute of Gravitation and Space Sciences

Gravitational Researches Laboratory

21 Ana Ipatescu bd., 71111 Bucharest, Romania

Abstract

The paths on the $\mathbf{R}^3$ real Euclidean manifold are defined as 2-dimensional simplicial strips. The Lagrangian of a moving mass is proportional to the width of the path. The special relativistic form of the Lagrangian is recovered in the continuum limit, without relativistic Lorenz invariance considerations¹.

1 Introduction

Standard approaches to discrete space-time consist in the description of the real space-time manifold as a geometrical lattice [1], [2], [3] or in the space-time approximation by a piecewise linear space [4], [5]. In this paper, we intend to introduce a method of discretization which is not performed upon the space-time manifold, but on the paths, which are defined as 2-dimensional simplicial strips embedded in an Euclidean space. Preliminary results were given in [6], [7].

One of the physical reasons in order to introduce the space-time discreteness is given by the experimental fact that the speed of light c is maximal and constant; it is easy to explain this by introducing space-time characteristic constants: an elementary length λ and an elementary time τ . If λ and τ are of Planck dimensions ($\lambda_P = (\hbar G/c^3)^{1/2} \simeq 1.616 \times 10^{-35}\text{m}$ and $\tau_P = \lambda_P/c \simeq 5.391 \times 10^{-44}\text{s}$) [8], it is generally accepted that no important influences occur on the continuum space-time concept at the usual scales

of physical phenomena. In the sequel, we consider $\lambda = \tau = 1$, if no other dimensional considerations are given.

Usual geometrical lattices do not present isotropy, i.e. some favoured directions are kept in the continuum limit. To circumvent this failure, we shall make use of a isotropic lattice, which covers the real manifold. Then, we introduce the simplicial paths. In the final part of the paper, a connection with the continuum special relativity is presented.

2 Simplicial paths

Let $\mathbf{R}^3$ the real linear space endowed with the Euclidean metric. It is almost obvious that if we accept an elementary length and we need rotational invariance, the elementary movements should cover the set of all the possible directions and senses in $\mathbf{R}^3$; this set is isomorphic to $S^2 \subset \mathbf{R}^3$ - the 2-sphere. In order to avoid confusions, we will denote by $\mathbf{x}$ the vectors of $\mathbf{R}^3$ and by x the points of the corresponding affine space. Let $|\mathbf{x}|$ the length of $\mathbf{x}$. Let $S^2(x_0)$ the unit length radius sphere with the center in $x_0 \in \mathbf{R}^3$. Let $\mathbf{Z}$ the ring of integers.

Theorem 1 *There exists at least one embedding of $\mathbf{R}^3$ in the free $\mathbf{Z}$ -module generated by $S^2(x_0)$. $\square$*

Proof. In order to prove the theorem, we show that each $\mathbf{x} \in \mathbf{R}^3$ has a realization in the $\mathbf{Z}$ -module. Let $a_i \in S^2(x_0)$. Let $\mathbf{a}_i \in \mathbf{R}^3$ the real unit length vectors with the directions given by a_j . An element $\bar{\mathbf{x}}$ of the free $\mathbf{Z}$ -module generated by $S^2(x_0)$ is identified with a real vector $\mathbf{x}$ as:

$$\bar{\mathbf{x}} = \sum_i z_i \mathbf{a}_i \equiv \mathbf{x}, \quad z_i \in \mathbf{Z} \quad (1)$$

This means that elements of $\mathbf{R}^3$ could be generated by combinations of unit length real translations $\mathbf{a}_i, |\mathbf{a}_i| = 1$, with integer coefficients. Let $\forall \mathbf{x} \in \mathbf{R}^3$; it is always possible to explicit the length of $\mathbf{x}$ as $|\mathbf{x}| = n + \epsilon$, where $n \in \mathbf{Z}^+$ (natural number) and $\epsilon \in [0, 1)$. Let $\mathbf{a}_x = \mathbf{x}/|\mathbf{x}|$ the corresponding unit vector of $S^2(x_0)$ possessing the same direction as $\mathbf{x}$. If $\mathbf{e} = \epsilon \cdot \mathbf{a}_x$, there always exist two unit vectors $\mathbf{a}_\epsilon$ and $\mathbf{a}'_\epsilon$ as $\mathbf{e} = \mathbf{a}_\epsilon + \mathbf{a}'_\epsilon$. In this way:

$$\mathbf{x} = n \cdot \mathbf{a}_x + \mathbf{e} = n \cdot \mathbf{a}_x + \mathbf{a}_\epsilon + \mathbf{a}'_\epsilon, \quad \forall \mathbf{x} \in \mathbf{R}^3 \quad \text{Q.E.D.} \quad (2)$$

Corollary 2 *Let $x, x' \in \mathbf{R}^3$. It is always possible to connect the points x and x' with at least one combination of $\mathbf{a}_i$ vectors of the form 1. $\square$*

Proof. Let $\mathbf{x} = \mathbf{vect}(x, x') \in \mathbf{R}^3$ the vector defined by the points x and x' , then see 2 Q.E.D.

Let us consider that a particle is confined in a space zone of Planck dimensions; it seems natural that the simplest geometrical model of the zone is the interior of a sphere $S^2(x)$ of Planck radius λ_P . If the particle is considered a point mass, the most probable movement due to symmetry considerations is the radial one: the particle "jumps" from the center of the sphere to the surface. If we take $\lambda_P = 1$, we may consider that the point particle "moves" acted by unit length translations $\mathbf{a}_i$; we will call the resultant trajectory a *d-path* (d from discrete).

Definition 1 *A d-path is the ordered set $d_{0N} = \{x_0, x_1, \dots, x_N\}$ of points as:*

$$\mathbf{vect}(x_0, x_1) = \mathbf{a}_1; \mathbf{vect}(x_1, x_2) = \mathbf{a}_2; \dots \mathbf{vect}(x_{N-1}, x_N) = \mathbf{a}_N, \mathbf{a}_k \in \mathbf{R}^3, |\mathbf{a}_k| = 1. \quad (3)$$

In order to describe a movement, we need some topological ingredients to get the continuity. For an example, if we join two-by-two the consecutive points x_k, x_{k+1} with segments, we get a 1-dimensional simplicial complex made of unit length edges. The polyhedra attached to such 1-dimensional simplicial complexes are called *edge-paths*; their properties are known in algebraic topology [9]. However, if we describe the trajectories by the mean of edge-paths, the immediate conclusion is that movement could develop only with the speed of light. In order to remove this difficulty, the following constructions are introduced:

Definition 2 *Let d a d-path defined by the set of points $\{x_0, x_1, \dots, x_N\}$. Let $a_k = \overline{x_{k-1}x_k}$ the edges defined by the points x_{k-1} and x_k . Let σ_k the 2-dimensional simplex (isosceles triangle) defined by the segments a_k and a_{k+1} . The **D-path** D is the polyhedron $|\Sigma|$ attached to the 2-dimensional simplicial complex Σ , as follows:*

$$\Sigma = \{x_0, x_1, \dots, x_N\} \cup \{a_1, a_2, \dots, a_k\} \cup \{\sigma_1, \sigma_2, \dots, \sigma_{k-1}\} \quad D \equiv |\Sigma|. \quad (4)$$

Definition 3 *The speed attached to the simplex σ_k component of D is the vector $\mathbf{v}_k$ defined by the baricentres $\hat{x}_{k-1}$ and $\hat{x}_k$ of the segments a_k and a_{k+1} which generate the simplex σ_k :*

$$\mathbf{v}_k = \mathbf{vect}(\hat{x}_{k-1}, \hat{x}_k) . \quad (5)$$

It is easy to note that $|\mathbf{v}_k| \in [0, 1]$ as in the special relativity theory. We mention that the procedure of baricentric subdivision is known in algebraic topology as a method to construct the derivative of a simplicial complex [9]; in this way, the edge $v_k = \overline{\hat{x}_{k-1}\hat{x}_k}$ of our construction may be regarded as the derivative of σ_k , or the derivative of D at time k .

The standard procedure to describe the movement of a point-like particle on the $\mathbf{R}^3$ manifold is to introduce the paths described by the particle, which are orbits of the action of some real one-parameter subgroup G on $\mathbf{R}^3$, in such a manner that:

$$[0, T] \xrightarrow{G} \mathbf{x}(t)|_0^T \quad (6)$$

with $\dot{\mathbf{x}}(t) \equiv d/dt[\mathbf{x}(t)]$ the speed. In the present approach, G is replaced by a discrete one-parameter subgroup $\mathcal{G}$, i.e.:

$$[0, 1, \dots, N] \xrightarrow{\mathcal{G}} \{x_0, x_1, \dots, x_N\} \xrightarrow{D-path} \{\sigma_1, \sigma_2, \dots, \sigma_{N-1}\} \quad (7)$$

(the lose of the end points of the discrete time interval is not significant for this approach). By means of the baricentric subdivision, we attached the speed $\mathbf{v}_k$ (see 5) to each simplex σ_k component of the $D-path$ $\{\sigma_1, \sigma_2, \dots, \sigma_{N-1}\}$ in such a manner that $\mathbf{v}(\sigma_k) = \mathbf{v}(k)$.

It is important to notice the difference between a simple discrete path, which is an ordered set of disconnected points of $\mathbf{R}^3$, and a $D-path$, which is an ordered set of geometric 2-simplexes connected by edges, whose topology is the one induced by the attached polyhedron (see [9]).

It is easy to observe the correspondence with the continuum mechanics: for large speed ($|v_k| \rightarrow 1$), the $D-path$ goes to a $d-path$, which is 1-dimensional; for small speed, the $D-path$ becomes a strip of width $\simeq \lambda$ and the speed goes to a vector tangent to the median line of the strip, line which identifies itself with the classical trajectory. In this sense, we shall define the continuum limit as follows.

Definition 4 Let D a D -path. The **1-dimensional path** attached to D is the polyhedron $|V|$ attached to the 1-dimensional simplicial complex V defined by the baricenters $\hat{x}_k$ of the edges of the corresponding d -path:

$$V = \{\hat{x}_0, \hat{x}_1 \dots \hat{x}_N\} \cup \{v_1, v_2, \dots v_{N-1}\}; \quad v_k = \overline{\hat{x}_{k-1} \hat{x}_k} \quad . \quad (8)$$

Within the upper definition, the 1-dimensional path is a triangulation [9] of a classical path, made of edges of length $l_k = |v_k| \in [0, 1]$ or, with dimensional considerations, $l_k \in [0, \lambda]$. The speed attached to the edge is $\mathbf{v}_k$ (see 5). The coordinate $\mathbf{x}_k$ of the edge may be considered the position of either $\hat{x}_k$, or the baricenter of v_k , or even the baricenter of σ_k . The time coordinate is $k \in \{0, 1 \dots N\}$ or, with dimensional considerations, $\tau \cdot k$. In this way, we may write:

$$\begin{aligned} \Delta \mathbf{x}(k) &= c \cdot \mathbf{v}_k \cdot \tau \quad \longrightarrow \quad d\mathbf{x} = \mathbf{v}(t) \cdot dt \\ \Delta l(k) &= c \cdot v_k \cdot \tau \quad \longrightarrow \quad ds = v(t) \cdot dt \end{aligned}$$

$$l_{0N} \equiv \text{the length of the path} = \lambda \cdot \sum_{k=1}^N v_k \quad \longrightarrow \quad \int_{x_0}^{x_N} ds = \int_{t_0}^{t_N} v(t) dt \quad (9)$$

We regain in this way the finite differences methods and go to the continuum limit by means of standard procedures.

3 The Lagrangian

From the preceding section, we reached to the concept of D -path, which contains by itself sufficient parameters in order to define the movement: *position* $\mathbf{x}_k$ and *speed* $\mathbf{v}_k$. We elude the concept of point mass, if we consider that the mass is distributed on the 2-simplex σ_k . In this case, in a good approximation, $\mathbf{x}_k$ may be identified with $\hat{\mathbf{x}}(\sigma_k)$ - the position of the baricenter of the simplex σ_k .

In order to construct the Lagrangian $L(\mathbf{x}_k, \mathbf{v}_k)$ for a mass m_0 moving on a D -path, we shall start from the following physical considerations:

- i) L must be invariant under the Euclidean rotations and translations, in order to preserve the fundamental properties of movement;
- ii) $L(\mathbf{x}_k, \mathbf{v}_k)$ must be a specific component of the D -path, which contains sufficient variables to describe the movement.

According to the geometric construction of the D -path, the magnitude v_k of the speed $\mathbf{v}_k$ on σ_k may be written as follows:

$$v_k = \cos \frac{\alpha_k}{2} \quad , \quad \alpha_k = \pi - \arccos(\mathbf{a}_{k-1} \cdot \mathbf{a}_k) \quad (10)$$

where α_k is the angle between the edges a_{k-1} and a_k which determine the simplex σ_k (see 4). In terms of speed v_k (see 8), the length of the height line (normal to v_k) of the geometrical simplex σ_k becomes:

$$h_k = \sin \frac{\alpha_k}{2} = \sqrt{1 - v_k^2} \quad (11)$$

proportional to the Lorentz square root.

In the continuum limit ($\lambda, \tau \rightarrow 0$ as $\lambda/\tau = c$), h_k identifies itself with the transversal width of the strip defined by the D -path at the position x_k . Using adequate units 9 (see also section I), we get for h_k :

$$\frac{h_k}{\lambda} = \sqrt{1 - \frac{v_k^2}{c^2}} \iff \frac{h(k)}{\lambda} = \sqrt{1 - \frac{v^2(k)}{c^2}} \xrightarrow{\text{continuum limit}} \sqrt{1 - \frac{v^2(t)}{c^2}} \quad (12)$$

The relativistic Lagrangian for a point particle of rest mass m_0 is defined in continuum relativistic mechanics as [8]:

$$L(v) = -m_0 c^2 \sqrt{1 - \frac{v^2(t)}{c^2}} \quad (13)$$

After dimensional considerations we get from 12 and 13 the following:

Definition 5 *The Lagrangian $L(v_k)$ on a D-path is defined as:*

$$L(v_k) \equiv -m_0 c^2 \frac{h_k}{\lambda} \quad (14)$$

proportional to the height h_k of the simplex σ_k .

In the continuum limit, we get:

$$L(v_k) = -m_0 c^2 \frac{h_k}{\lambda} \longrightarrow -m_0 c^2 \frac{h(t)}{\lambda} = -m_0 c^2 \sqrt{1 - \frac{v^2(t)}{c^2}} \quad (15)$$

that the simplicial Lagrangian 14 goes to the standard relativistic Lagrangian. This is surprising, because no relativistic invariant has been involved in the construction of the model. It is also interesting to note that the Lagrangian 14 is independent on the magnitude of the elementary length λ .

4 Conclusions

The starting point of our analysis is the idea that the speed of light is maximal and constant due to some discrete properties of the space-time. In the present approach, we did not discretize the space-time manifold, but the paths. First, we took as base manifold the $\mathbf{R}^3$ Euclidean space and we considered that an elementary discrete movement is a set of sequentially equidistant points of $\mathbf{R}^3$. In order to emphasize the rotational invariance, we showed in the second section that these sets are elements of the free $\mathbf{Z}$ -module generated by the $S^2 \subset \mathbf{R}^3$ unit radius sphere and we proved that all $\mathbf{R}^3$ is covered by means of this procedure. In order to give a topology (to introduce the continuity) on the discrete sets of points, we defined the *D-paths* as 2-dimensional simplicial complexes having as vertices these sets. By means of a baricentric subdivision, we attached to each simplex component of the D-path a *speed* whose magnitude varies between 0 and 1. In the third section, we defined the Lagrangian of a moving mass (distributed on a 2-simplex) as being proportional to the transversal width of the simplicial strip. In the continuum limit, the Lagrangian becomes the correspondent relativistic one for the point particle. In this way, we are able to claim that the basic special relativistic properties of movement may be derived as a consequence of the local discreteness of the space-time manifold.

References

- [1] H. Yamamoto, Phys. Rev. **D 30**, 1727 (1984), **D 32**, 2659 (1985)
- [2] K.I. Macrae, Phys. Rev.**D 23**, 886 (1981)
- [3] M. Lehto, H.B. Nielsen, M. Ninomyia, Nucl. Phys. **B 272**, 213,228 (1986).
- [4] T. Regge, Nuovo Cimento **XIX**, 558 (1961)
- [5] R.M. Williams and P.A. Tuckey, Class. Quantum Grav. **9**, 1409 (1992).
- [6] M.I. Piso, Nuovo Cimento **108** B, no 11, 1307 (1993)
- [7] M.I. Piso, S. Sainar, UTFT preprint 9|92 (1992).
- [8] C.W. Misner, K.S. Thorne, J.A. Wheeler, *Gravitation* (W.H. Freeman, 1973).
- [9] C.R.F. Maunder, *Algebraic Topology* (Cambridge University Press, 1980).

¹This work has been partially supported on the Romanian Space Agency 1013C contract.